

Thursday Sep 25th

→ SIT WITH YOUR GROUP TODAY! ←

Topic # 6 Atmospheric Structure & Chemical Composition

- **Self Test 4 & RQ-4 on The Laws of Thermodynamics** are now posted. The readings that will prepare you for this Topic's Self Test & RQ are listed in Self Test 4 and next week's CHECKLIST.
- RQ-4 is **DUE NEXT TUESDAY 30** minutes before class!
- Test #1 grade worries? Looking for more bonus point chances? Details **TODAY in class** on another one!
- The **LINKING-TO-LIFE PROJECT** directions will be posted this weekend. **Read through them before class next Tuesday** and come in with your questions.

◆ **REMINDER:**


It's that time during the semester when requests for **Grade Reports** start coming up . . .

SEE FAQ # 29

<http://fp.arizona.edu/kkh/nats101gc/faq.htm#29>

If you need me to sign a grade report -- notify me in advance by email at least one day before you need me to sign your grade report!!

ANSWERS to the last part of G-1:



All 3 are illustrating ABSORPTION of incoming Solar SW by the EARTH's surface followed by outgoing RADIATION of LW Infrared from the EARTH's surface !

Q7. Which diagram above shows SW (solar radiation being reflected back to space?


A

B


C

None of them

p 33



Here's the correct diagram to show **SW** Solar being reflected back to space!



Q8. Diagram A shows LW (IR) terrestrial radiation “**bouncing off**” (or **reflecting**) the gases in the atmosphere and being sent back to Earth’s surface. (i.e. being reflected back to the surface by the gases without being absorbed by them.)

Is this an accurate depiction of how the Greenhouse Effect works?


Yes

No

Why or Why not?

The LW (IR) radiation is **absorbed** by the GHG’s and then **re-radiated (or re-emitted)** back down to the Earth’s surface to warm it. The IR is **NOT reflected**.
IF IT’S REFLECTED IS NOT ABSORBED.

DON’T USE “BOUNCING or “REFLECTING” to describe the Greenhouse Effect process: **GH gases ABSORB & RE-RADIATE!**



Q9. Diagram B shows LW (IR) terrestrial radiation being **absorbed and then emitted back down** by the gases in the atmosphere.

Is this an accurate depiction of how the Greenhouse Effect works?


Yes


Why or Why not?

(although it's not quite complete – more on this later)

LW (IR) being absorbed by GH Gases, and then emitted out again, (back down to the surface of the Earth) is **exactly how the GH Effect operates.**

Example: CO₂ absorbs IR wavelengths and then emits IR wavelengths in this familiar animation:





Q10. Diagram C shows LW (IR) terrestrial radiation **going right through the atmosphere** out to space.


Is this an accurate depiction of how the Greenhouse Effect works?

Yes No Why or Why not?

Diagram C shows **ALL** the IR **leaving the Earth's surface** and **NOT** being absorbed at all !

If this happened the Earth's temperature would be below freezing!

Q11. On the diagram that you think **best depicts the processes involved in the GREENHOUSE EFFECT**, CIRCLE the specific part of the diagram that represents the **Greenhouse Effect**:



DO NOT CIRCLE any part of the **SUN's incoming SW!**
(Shortwave Radiation = IR + VIS)
SW is NOT part of the GH Effect!




THINKING MORE DEEPLY . . .

① Some Incoming SW radiation from the SUN goes right through the atmosphere to Earth (w/o being absorbed)


How? Through the UV-VIS Atmospheric Window!

③ Some IR radiation is emitted from the Earth's surface right out to space through "IR window"

Outgoing LW



⑤ Some IR radiation is **absorbed** by GH gases in the atmosphere, but is **emitted out to space** (not back to Earth)



④ Some IR radiation is **absorbed** by GH gases in the atmosphere and **emitted back to Earth**

② The Earth absorbs SW that reaches the surface

Absorption & re-emission by GH gases


Absorption & re-emission by GH gases

Bottom of p 33

There's one more thing to correct in our the depiction of incoming Solar

Some SW radiation
gets absorbed on
its way down to the
surface!

(in addition to
terrestrial LW (IR)
radiation being
absorbed in the GHE)



**Topic # 6 ATMOSPHERIC
STRUCTURE
&
CHEMICAL COMPOSITION**

**All about the GASES IN THE
ATMOSPHERE, esp.
GREENHOUSE GASES!**

Class Notes pp 35- 39


OBJECTIVES:

To understand:




- the **VERTICALSTRUCTURE** of the atmosphere & its relationship to temperature
- which **GASES** are in the atmosphere
- **where** they are concentrated, and
- why gases at different levels are linked to the **Greenhouse Effect & Ozone Depletion**







Things you've seen before that will all come together under this topic:



$$E = \sigma T^4$$



$$E = hc / \lambda$$




$$\lambda_m = a / T$$

**We travel together, passengers in a
little space-ship, dependent on its
vulnerable supplies of air and soil.**

~ Adlai Stevenson




The atmosphere has a “structure” of different named layers :



These layers have different thicknesses and temperatures...

Most everything WE experience is in the lowest layer, the **TROPOSPHERE**

This zig-zag line is showing changes in temperature with altitude



The Vertical Structure of the Atmosphere

KEY CONCEPT:

The atmosphere's vertical structure is defined by **CHANGES** in the trend of **TEMPERATURE** with height.






Figure 3-9b in SGC E-text

Atmospheric Pressure = weight of the air column above

Atmospheric Pressure & Mass Vary with Height




99% of mass lies below ~ 50 km (top of Stratosphere)

50% of mass lies below ~ 6 km (middle Troposphere)




The changes in temperature with height are the result of: differential absorption of shortwave (SW) & longwave (LW) radiation by atmospheric GASES concentrated at various altitudes.




Here's why
these
changes in
temperature
occur :

Let's start at the
SURFACE →




A nice online review . . .

<http://earthguide.ucsd.edu/earthguide/diagrams/atmosphere/index.html>



REVIEW: The pattern of electromagnetic wavelengths that are **absorbed & emitted** by a particular atom (or combination of atoms)


is called its **ABSORPTION SPECTRUM** or its **ABSORPTION CURVE**



**The Absorption curve for
Ozone / Oxygen**



How incoming **SOLAR** radiation of different wavelengths gets **TRANSMITTED** or **ABSORBED** by different gases on its way to the Earth's surface



1. UV, $\lambda < 0.12 \mu\text{m}$, absorbed by N₂ and O₂ in upper atmosphere
2. UV, $0.12 \mu\text{m} \leq \lambda < 0.18 \mu\text{m}$ absorbed by O₂
3. UV, $0.18 \mu\text{m} \leq \lambda < 0.34 \mu\text{m}$ absorbed by O₃ in ozone layer
4. Near UV and visible, $0.34 \mu\text{m} \leq \lambda < 0.7 \mu\text{m}$ transmitted nearly undiminished except for scattering
5. Near IR, $0.7 \mu\text{m} \leq \lambda < 3.0 \mu\text{m}$, absorbed slightly by O₂ and in troposphere by H₂O

Reminder: Ultraviolet radiation: UVC = 0.20 - 0.29 UVB = 0.29 - 0.32 UVA = 0.32 - 0.40 μm


GROUP WORK:
Study this box
of info & answer
Q1, Q2, & Q3

**AFTER YOU'VE
WORKED ON
page 36
in YOUR GROUP . . .**

**CLICKER
your answers in!**


Clicker Q 1. The **GREATEST** amount of incoming solar energy (represented by the width of the arrows) is transferred to Earth via **which wavelengths** of electromagnetic radiation?

1. UV $< 0.12 \mu\text{m}$
2. UV $0.12 - 0.18 \mu\text{m}$
3. UVC + UVB
4. BOTH arrows
- 4 + 5




Clicker Q 1. The **GREATEST** amount of incoming solar energy (represented by the width of the arrows) is transferred to Earth via **which wavelengths** of electromagnetic radiation?

1. UV $< 0.12 \mu\text{m}$
2. UV $0.12 - 0.18 \mu\text{m}$
3. UVC + UVB
4. BOTH arrows 4 + 5



Clicker Q 2. Why does ARROW #3's radiation get attenuated below 50 km?

1. Because this is the area of the **mesosphere** and there is very little absorption of radiation in this layer
2. Because **nitrogen (N₂)** and **oxygen (O₂)** are abundant at 50 km and act as GHG's to **absorb** the **UVC + UVB** rays
3. Because this is the area of the stratosphere where **ozone (O₃)** is **concentrated** and absorbs harmful **UVC + UVB** rays




Clicker Q 2. Why does ARROW #3's radiation get attenuated below 50 km?

1. Because this is the area of the **mesosphere** and there is very little absorption of radiation in this layer
2. Because **nitrogen (N₂)** and **oxygen (O₂)** are abundant at 50 km and act as GHG's to **absorb** the **UVC + UVB** rays
3. Because this is the area of the stratosphere where **ozone (O₃)** is **concentrated** and absorbs harmful **UVC + UVB** rays




Clicker Q 3. Why does ARROW #5's radiation get attenuated (thinner) below 10 km?

1. Because **ozone (O_3)** is abundant below 10 km and absorbs large amounts of incoming **IR**
2. Because this is the area of the troposphere where **water vapor (H_2O)** is abundant and (as a GHG) it **absorbs IR**
3. Because **clouds** in the troposphere block out some of the incoming **visible light** rays



Clicker Q 3. Why does ARROW #5's radiation get attenuated (thinner) below 10 km?


1. Because **ozone (O_3)** is abundant below 10 km and absorbs large amounts of incoming **IR**
2. Because this is the area of the troposphere where **water vapor (H_2O)** is abundant and (as a GHG) it **absorbs IR**
3. Because **clouds** in the troposphere block out some of the incoming **visible light** rays



UV rays < .32 μm
very harmful to
life on Earth arrows
1, 2 + 3




How incoming
SOLAR
radiation of
different
wavelengths
gets
TRANSMITTED
or ABSORBED
by different
gases
on its way to
the Earth's
surface



1. UV, $\lambda < 0.12 \mu\text{m}$, absorbed by N₂ and O₂ in upper atmosphere
2. UV, $0.12 \mu\text{m} \leq \lambda < 0.18 \mu\text{m}$ absorbed by O₂
3. UV, $0.18 \mu\text{m} \leq \lambda < 0.34 \mu\text{m}$ absorbed by O₃ in ozone layer
4. Near UV and visible, $0.34 \mu\text{m} \leq \lambda < 0.7 \mu\text{m}$ transmitted nearly undiminished except for scattering
5. Near IR, $0.7 \mu\text{m} \leq \lambda < 3.0 \mu\text{m}$, absorbed slightly by O₂ and in troposphere by H₂O


Reminder: Ultraviolet radiation: UVC = 0.20 - 0.29 UVB = 0.29 - 0.32 UVA = 0.32 - 0.40 μm

Incoming SW




OK – so that explains what happens in different layers of the atmosphere to the **INCOMING SOLAR Shortwave (SW)** on its way down to the Earth's surface

Outgoing LW



. . . But what happens to the **OUTGOING TERRESTRIAL Longwave (IR)** radiation when it radiates from the Earth's surface upwards??

Write in the names of the layers:



EXPLORING THE
EVIDENCE . . .



**The Greenhouse
Warming Signature:**
*"Increasing CO₂ warms
the Troposphere and
cools the Stratosphere"*

The "Greenhouse Effect" Warming Signature

The Greenhouse Signature



What would a SOLAR Warming Signature look like?

ATMOSPHERIC COMPOSITION

Which gases?

What concentration?

Which ones are

Greenhouse Gases (GHG)?

Where do the GHG's come from?

**Which GHG's are changing in
concentration due to**

HUMAN ACTIVITIES?



ATMOSPHERIC COMPOSITION



* = Greenhouse Gas (GHG)

RF = Radiative Forcing of GHG's in Wm^{-1}

Gas	Symbol	Percent Concentration (by volume dry air)	Concentration in Parts per Million (ppm)	*RF W/m^2
Nitrogen	N ₂	78.08	780,800	
Oxygen	O ₂	20.95	209,500	
Argon	Ar	0.93	9,300	
* Water Vapor	H ₂ O	0.00001 (<i>South Pole</i>) – 4 (<i>Tropics</i>)	0.1 (<i>South Pole</i>) – 40,000 (<i>Tropics</i>)	<i>varies</i>
* Carbon Dioxide	CO ₂	0.0390+ (2009) http://co2now.org/	390+ (2010) http://co2now.org/	1.66
* Methane	CH ₄	0.0001774 (<i>in 2005</i>)	1.774	0.48
* Nitrous Oxide	N ₂ O	0.0000319	0.319	0.16
* Ozone	O ₃	0.0000004 (<i>in 70s</i>)	0.01 (<i>at the surface</i>)	<i>varies</i>
* CFCs (e.g. Freon-12) (Chlorofluorocarbons)	CCl ₂ F ₂	0.0000000538	0.000538 <i>RF for all CFC Totals:</i>	0.170 0.268
* HCFCs (e.g., HCFC-22) (Hydrochlorofluorocarbons)	CHClF ₂	0.0000000169	0.000169 <i>RF for all HCFC Totals:</i>	0.033 0.039
Neon, Helium, Hydrogen, Krypton, Xenon	Ne, He, H, Kr, Xe	0.0018 – 0.000009	18 – 0.09	
Particles (dust, soot)	--	0.000001	0.0001	



For more on GHG concentrations see: <http://www.ipcc.ch/pdf/assessment-report/ar4/wg1/ar4-wg1-chapter2.pdf> Table 2.1

Most Abundant Gases in the Atmosphere

GAS	Symbol	% by volume	% in ppm
Nitrogen 	N₂	78.08	780,000
Oxygen 	O₂	20.95	209,500
Argon	Ar	0.93	9,300

↓
Total = 99.96%

Next Most Abundant Gases:

GAS	Sym bol	% by volume	% in ppm
<p>Water Vapor</p>  <p>The diagram shows two water molecules (H₂O) with a central carbon atom (C) and two hydrogen atoms (H). The first molecule is labeled 'Slow rotation rate' and the second is labeled 'Faster rotation rate'. An arrow points from the first to the second, indicating a transition or change in rotation rate.</p>	<p>H₂O</p>	<p>0.00001 (South Pole) to 4.0 (Tropics)</p>	<p>0.1 - 40,000</p>
<p>Carbon Dioxide</p>  <p>The diagram shows a carbon dioxide molecule (CO₂) with a central carbon atom (C) and two oxygen atoms (O). The molecule is shown in a bent configuration, with arrows indicating the bending motion. The label 'Bending mode' is placed below the diagram.</p>	<p>CO₂</p>	<p>0.0390 (and rising!)</p>	<p>360 (in 1997) 390 ! (in May 2009)</p>

Greenhouse Gases !

Other Important Greenhouse Gases:


GAS	Symbol	% by volume	% in ppm
Methane	CH₄	0.00017	1.7
Nitrous Oxide	N₂O	0.00003	0.3
Ozone	O₃	0.00000004	0.01
CFCs (Freon-11)	CCl₃F	0.0000000026	0.00026
CFCs (Freon-12)	CCl₂F₂	0.0000000047	0.00047

Greenhouse Gases!

CO₂

Amount in
Atmosphere = 400,000+ ppb

(From: DP text p 29 where it says 386,000 ppb!)



With your Group . . .
STUDY THE TABLE ON

Page 38

to familiarize yourself with
each of the GHG's

Then get ready for the
“NAME THAT GAS!”

TEAM COMPETITION




Infrared Radiators

GROUPS: # 1, # 2, #3, #4


Name that GAS!!!

MYSTERY

GHG # 1



METHANE: Trends



METHANE (CH₄): Sources

* Produced naturally in anaerobic processes (e.g., decomposition of plant material in swamps & bogs)

* **Has increased** due to the following activities: **raising cattle / livestock, rice production, landfill decomposition, pipeline leaks**

* **Has relatively short atmospheric residence time** because it reacts with OH (~10 years)




Tangerine Tazers
GROUPS: # 5, 6, 7, 8

Name that **GAS!!!**

MYSTERY
GHG #2



CARBON DIOXIDE: Trends



CARBON DIOXIDE :

* Arrives in atmosphere naturally through the natural carbon cycle

* Has increased dramatically since the 1800s due to:

FOSSIL FUEL COMBUSTION:

oil, coal, gas (automobiles) . . .

But especially **COAL**

CARBON DIOXIDE (cont.):

* **RESIDENCE TIME** in the atmosphere of **CARBON ATOMS** in the carbon cycle = **~ 12.7 years;**

but **residence time of CO₂ GAS MOLECULES** is estimated at about 100 years

Plus it takes **50 to 100 years** for atmospheric **CO₂ to adjust** to changes in sources or sinks.

If we make changes now, it will still be many, many years before the effect will be felt!



Mellow Yellow Reflectors
GROUPS: # 9, 10, 11, 12


Name that GAS!!!

MYSTERY

GHG # 3



NITROUS OXIDE: Trends



NITROUS OXIDE (N₂O): Sources



- * Produced naturally in soils
- * Has increased due to fossil fuel combustion (esp. diesel), forest burning, use of nitrogen fertilizers
- * Has long atmospheric residence time (~ 150 years)




Greenhouse Gassers
GROUPS #13,14, 15, 16

Name that GAS!!!



MYSTERY
GHG # 4



CFCs: Trends



Human-made --
didn't exist
before 1950!



CFCs (Freon-11 & Freon-12)

- * Human-made CFCs (didn't exist in atmosphere prior to 1950s)

- * Have increased at rates faster than any other greenhouse gas; used in refrigerants, fire retardants, some aerosol propellants & foam blowing agents

- * Absorb at different wavelengths than H₂O and CO₂ (in 8 –12 μm “WINDOW” part of spectrum), hence a single molecule can have great effect

MONTREAL (and subsequent) PROTOCOLS have reduced CFCs!

Clicker Q 4 – Why do you think the concentration of CFC's didn't begin dropping immediately after the Montreal Protocol in 1987?

- 1. Because it was an international “agreement only” and the nations of the world never followed through.**
- 2. Because it called for only a 50% reduction of CFC's over 10 years and had to be followed by more stringent protocols later.**
- 3. Because CFC's are very stable molecules and don't break down easily once they are in the atmosphere.**

Clicker Q4 – Why do you think the concentration of CFC's didn't begin dropping immediately after the Montreal Protocol in 1987?

- 1. Because it was an international “agreement only” and the nations of the world never followed through.**
- 2. Because it called for only a 50% reduction of CFC's over 10 years and had to be followed by more stringent protocols later.**
- 3. Because CFC's are very stable molecules and don't break down easily once they are in the atmosphere.**


Blue Sky Diffusers

GROUPS: # 17, 18, 19, 20

Name that **GAS!!!**


MYSTERY GHG # 6






WATER VAPOR

- * Arrives in atmosphere naturally through evaporation & transpiration
- * Due to unique quantum rotation frequency, H₂O molecules are excellent absorbers of IR wavelengths of **12 μ m and longer**;



GAS Table on
p 38

Virtually 100% of IR longer than 12 μm is absorbed by H_2O vapor and CO_2



Graphs on p 38 Class Notes

(12 μm close to the radiation wavelength of 10 μm , at which most of Earth's terrestrial radiation is emitted.)

IR at 12 μm absorbed




WATER VAPOR (cont):

* H₂O has variable concentration and residence time in the atmosphere depending on location and atmospheric circulation

Blue = wettest climates, lots of humidity & water vapor


Yellow = driest climates, less atmospheric water vapor





At higher air temperatures, H₂O molecules collide & rebound more frequently, leading to expansion of the air & the water vapor in the air.



Hence hot climates can hold more water vapor in the air



At lower air temperatures as air gets more dense, H₂O molecules are more likely to bond so that a phase change to liquid water or even solid ice can occur.



Hence in cooler climates, more of the available H₂O is likely to be in the liquid or solid state on the Earth's surface



WATER VAPOR (cont):

* H₂O is **NOT** globally increasing in direct response to human-induced factors, but if global temperatures get warmer, H₂O vapor in the atmosphere will increase

Why???

. . . due to more evaporation
in the warmer climate!

THINK ABOUT THIS!




OPEN FLOOR



Name that **GAS!!!**

(this one's a visual hint only!)

MYSTERY GHG # 7



OZONE: Sources





Produced naturally in photochemical reactions in STRATOSPHERIC ozone layer -- “good ozone”



Has increased in TROPOSPHERE due to photochemical smog reactions -- “bad ozone”

O₃ absorbs IR radiation of 9.6 μm, close to wavelength of maximum terrestrial radiation (10 μm)



**WHOLE
ATMOSPHERE**

Review)



OZONE:

Trends

Stratospheric ozone varies by latitude and season -- is affected by **solar radiation**, **volcanic eruptions** & **chemical reactions** due to CFCs.

Overall, O₃ is **decreasing** in the STRATOSPHERE

*More on OZONE later
on in the semester*





OPEN FLOOR

Name that

GAS!!!


This one's a repeat of a previously
guessed gas!

MYSTERY

GHG #5



CARBON DIOXIDE --- Trends:



The Keeling Curve




CARBON DIOXIDE (cont.):

*** Has increased dramatically since the 1800s due to:**


DEFORESTATION -- which has the effect of increasing the amount of carbon in the atmospheric “reservoir” by reducing the photosynthesis outflow and increasing the respiration inflow.

(Deforestation also accelerates forest decomposition, burning, etc. adding to the overall respiration inflow.)




CARBON emissions into the atmosphere are increasing:




Concentrations of Greenhouse Gases from 0 to 2005



Updated figures from **Dire** **Predictions** p 33





See you **MONDAY** at 5:30 pm
at Dr. Yoram Bauman's Presentation

ARIZONA  **WILDCATS**

GO CATS!
Beat the Ducks